



PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Bc

Search PubMed

for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display

Abstract

Show:

20

Sort

Send to

Text

Text Version

☐ 1: J Immunol Methods 1995 Nov 16;187(1):23-31

Links

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

**ELSEVIER SCIENCE
FULL-TEXT ARTICLE****Hydrogen peroxide destaining: a new method for removing non-specific stains in nitrocellulose membrane-based dot-ELISA for the detection of trypanosomes in tsetse flies (*Glossina* spp.).****Bosompem KM, Assoku RK, Nantulya VM.**

Noguchi Memorial Institute for Medical Research (NMIMR), Accra, Ghana.

Gut samples prepared from laboratory-reared tsetse flies and applied in dots onto nitrocellulose (NC) membrane were found to stain the membrane with differing coloration and intensity. The stains were, predominantly, either reddish to brown or blackish-brown to black and occasionally greenish to almost colourless, depending on the stage of digestion of the bloodmeal in the fly. NC membrane strips applied with tsetse gut samples from *T. brucei* infected and uninfected control flies were tested with the standard antigen detection dot enzyme-linked immunoassay (dot-ELISA), using a *T. brucei* specific monoclonal antibody (MoAb) and horseradish peroxidase goat anti-mouse conjugate. The stains in both infected and uninfected sample dots persisted through the assay. Furthermore, the staining intensity of some assayed uninfected sample dots were enhanced as a result of non-specific reactivity, making it difficult to distinguish between the infected and uninfected flies. This necessitated the development of a simple technique by which the non-specific stains and reactions could be removed. Sample 'dotted' NC membrane strips were destained by incubation with 5% hydrogen peroxide (H₂O₂) diluted in 5% skimmed milk in Tris buffer, pH 8.0. After washing, the destained strips were tested in the dot-ELISA. This method gave satisfactory reproducible results, since the most intense stains could be removed, and it had no effect on trypanosome antigens detected by a panel of four *T. brucei* species-specific, three *T. vivax* species-specific, four *T. congolense* species-specific and four *Nannomonas* subgenus-specific MoAbs. Using the destaining process in a modified dot-ELISA, 86 out of 95 (90.5%) of *Glossina morsitans centralis* flies experimentally infected with *T. brucei*, were identified. The destaining method was also used successfully to decolorize NC membrane bound tsetse faecal material.

PMID: 7490455 [PubMed - indexed for MEDLINE]

Display Show: Sort Send to

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer

May 20 2003 11:14:45